

Current Biology

Genetic and Epigenetic Strategies Potentiate Gal4 Activation to Enhance Fitness in Recently Diverged Yeast Species

Highlights

- *GAL* epigenetic transcriptional memory promotes faster reactivation of *GAL* genes
- Previously produced Gal1 provides a large future fitness benefit in galactose
- Epigenetic *GAL* memory most likely evolved from genetically encoded priming
- The Gal4 central domain promotes both Gal80 repression and maximal activation

Authors

Varun Sood, Jason H. Brickner

Correspondence

j-brickner@northwestern.edu

In Brief

Sood and Brickner find that faster *GAL* gene induction during transcriptional memory accelerates adaptation to galactose and requires both previously produced Gal1 to relieve Gal80 repression and the Gal4 central domain, which potentiates stronger activation. Related yeasts use either genetic or epigenetic means to alter *GAL* gene activation rates.



Genetic and Epigenetic Strategies Potentiate Gal4 Activation to Enhance Fitness in Recently Diverged Yeast Species

Varun Sood¹ and Jason H. Brickner^{1,2,*}¹Department of Molecular Biosciences, Northwestern University, Evanston, IL 60208, USA²Lead Contact*Correspondence: j-brickner@northwestern.edu<https://doi.org/10.1016/j.cub.2017.10.035>

SUMMARY

Certain genes show more rapid reactivation for several generations following repression, a conserved phenomenon called epigenetic transcriptional memory. Following previous growth in galactose, *GAL* gene transcriptional memory confers a strong fitness benefit in *Saccharomyces cerevisiae* adapting to growth in galactose for up to 8 generations. A genetic screen for mutants defective for *GAL* gene memory revealed new insights into the molecular mechanism, adaptive consequences, and evolutionary history of memory. A point mutation in the Gal1 co-activator that disrupts the interaction with the Gal80 inhibitor specifically and completely disrupted memory. This mutation confirms that cytoplasmically inherited Gal1 produced during previous growth in galactose directly interferes with Gal80 repression to promote faster induction of *GAL* genes. This mitotically heritable mode of regulation is recently evolved; in a diverged *Saccharomyces* species, *GAL* genes show constitutively faster activation due to genetically encoded basal expression of Gal1. Thus, recently diverged species utilize either epigenetic or genetic strategies to regulate the same molecular mechanism. The screen also revealed that the central domain of the Gal4 transcription factor both regulates the stochasticity of *GAL* gene expression and potentiates stronger *GAL* gene activation in the presence of Gal1. The central domain is critical for *GAL* gene transcriptional memory; Gal4 lacking the central domain fails to potentiate *GAL* gene expression and is unresponsive to previous Gal1 expression.

INTRODUCTION

Transcriptional adaptation to fluctuations in nutrient availability is critical for fitness [1]. In response to previous experiences, certain inducible genes show a mitotically heritable increase in the rate of transcription [2–12]. This epigenetic phenomenon, referred as transcriptional memory, is observed in yeast, *Drosophila*, and humans [5, 7, 13]. However, its evolutionary

history and adaptive impact have not been explored. Also, whereas some aspects of transcriptional memory are deeply conserved, gene-specific features also occur [4, 6, 8, 14], suggesting that gene-specific regulatory systems can be regulated by transcriptional memory.

In *S. cerevisiae*, *GAL* genes exhibit transcriptional memory. These genes are specifically induced in galactose to mediate galactose utilization [15, 16]. When cells are shifted from glucose to galactose, the initial rate of induction of *GAL* genes is very slow; the Gal1 protein reaches steady-state levels ~10 hr after shifting from glucose to galactose (Figure 1B; compared with the doubling time of budding yeast, 1.5–2 hr). However, in cells that have been previously grown in galactose, *GAL* genes are rapidly reactivated. This phenomenon persists in the population for 7 or 8 cell divisions after repression in glucose [2, 11, 14, 17]. Like other genes that show memory, *GAL* transcriptional memory is also associated with changes in chromatin structure, leading to a state that is poised for faster reactivation [14].

The initial induction of *GAL* genes is slow in part because, in most cells, there are fewer molecules of the Gal3 co-activator than the Gal80 inhibitor [11, 18]. Consequently, the expression is initially heterogeneous, with some cells responding and others not [18]. On the other hand, during memory, reactivation is more uniform [18]. *GAL* memory requires the Gal1 galactokinase, a paralog of Gal3 [2, 11, 14, 17]. Gal1 is abundantly produced in galactose, extremely stable, and diluted very slowly after repression in glucose [14]. A few hundred molecules of Gal1 are necessary and sufficient to produce the effects of memory, including the changes in chromatin structure over the *GAL1* promoter and uniformity of expression [11, 14]. Because both Gal1 and Gal3 can interact with Gal80 to relieve inhibition of the Gal4 activator [19–23], this leads to a model whereby residual Gal1 allows cells to more rapidly overcome Gal80 repression and rapidly induce *GAL* genes during memory (Figure 1A).

Here, we explored the adaptive value, evolutionary history, and molecular mechanism of *GAL* gene transcriptional memory. In *S. cerevisiae*, *GAL* transcriptional memory confers a strong adaptive advantage, allowing much faster adaptation to galactose and better utilization of mixed sugars. In contrast, *S. uvarum*, a divergent *Saccharomyces* species, does not benefit from previous growth in galactose but instead shows constitutively fast *GAL* gene activation due to basal *GAL1* expression. Replacing the *GAL1* promoter in *S. cerevisiae* with that from *S. uvarum* recapitulated this difference in behavior, confirming that the basal expression of Gal1 and the resulting fast activation




Figure 1. Expression Changes during GAL Memory and Their Fitness Effect

For a Figure360 author presentation of Figure 1, see the figure legend at <https://doi.org/10.1016/j.cub.2017.10.035>.

(A) Model for *GAL1* regulation and memory. Top: during activation, Gal3 sequesters the Gal80 repressor from the Gal4 activator, leading to *GAL* gene expression. Bottom: during reactivation, residual Gal1 augments Gal3 co-activation, leading to faster expression kinetics.

(B–G) Naive cells (ACT), naive cells expressing ectopic Gal1 (ACT + eGAL1), or cells that were grown in galactose overnight and shifted to glucose for 12 hr (REACT) were shifted to media containing galactose and either *GAL1*-mCherry fluorescence (B–D and F) or optical density 600 (OD₆₀₀) (E and G) were measured. *GAL1*-mCherry fluorescence was measured relative to constitutively expressed CFP using flow cytometry. (C and D) Effect of Gal80 inhibition on Gal1-mCherry expression. (D) Overlay of histograms for Gal1-mCherry from corresponding strains at the indicated times in (B) and (C). (E) At time = 0, all cultures were diluted to an OD₆₀₀ of 0.1 in galactose, and OD₆₀₀ was measured every 20 min using 96-well plate reader. Open circles represent the ratio of OD₆₀₀ between REACT and ACT. (F) Gal1-mCherry levels relative to CFP control after 7 hr in different concentration of galactose, plotted as fraction of expression in 1% galactose. (G) Growth and Gal1-mCherry expression (inset) in 0.2% glucose + 1.8% galactose. **p* ≤ 0.05; Student's *t* test.

Error bars for Gal1-mCherry fluorescence represent SEM from ≥3 biological replicates. The line and the bounding envelope for the OD₆₀₀ measurements are the mean and SEM, respectively, from ≥5 biological replicates. The yeast strains and the number of biological replicates for all experiments are listed in Tables S1 and S2, respectively.

of other *GAL* genes are genetically encoded by this promoter. Using a point mutation in Gal1 that disrupts the interaction with Gal80, we demonstrate that fast induction of *GAL* genes in both species results from a physical interaction between Gal1 and Gal80. A screen for mutants that block *GAL* memory downstream of Gal1 identified a mutation in Gal4 central domain (CD), revealing a critical role for this domain in transcriptional memory downstream of both Gal1 and Gal80. The CD serves to potentiate Gal4 activation, and this function is regulated by Gal80. Ectopically expressed CD complemented deletion of the CD in *trans* and was recruited to chromatin-bound Gal4Δcd, suggesting that potentiation is the result of an inter-domain interaction within Gal4. Thus, recently diverged species employ either

epigenetic or genetic strategies to alter the transcriptional potency of a transcription factor, promoting faster adaptation to changes in carbon source.

RESULTS

Transcriptional Memory Enhances Fitness by Promoting Uniform, Rapid Activation of *GAL* Genes

Because Gal1 is both necessary and sufficient to promote faster induction of *GAL* genes during memory, Gal1 most likely interacts with Gal80 to allow rapid de-repression of *GAL* genes (Figure 1A). The relative rates of *GAL1* transcription can be compared by measuring Gal1-mCherry fluorescence expressed




Figure 2. Genetic Screen for Mutants Defective for GAL Memory Identifies *gal1-D117V*

(A) Schematic of the 2-step FACS-based screen (see STAR Methods for details).

(B) Gal1-mCherry intensity relative to CFP internal control in wild-type and *gal1-D117V* mutant, measured by flow cytometry. Cells were shifted from glucose to galactose for activation (ACT) or grown in galactose overnight, shifted to glucose for 12 hr, and then shifted to galactose for reactivation (REACT). Error bar represents SEM from ≥ 4 biological replicates.

(C) Growth of wild-type and *gal1-D117V* mutant cells assayed by measuring OD₆₀₀ every 20 min during continuous growth in galactose (gal \rightarrow gal), during activation (ACT) or reactivation (REACT) after 12 hr of repression. The line represents the mean, and the envelope represents the SEM from ≥ 4 biological replicates.

(D) Co-crystal structure between Gal3 (pink) and Gal80 (blue), highlighting the salt bridge between the Gal3-Asp111 and Gal80-Arg367 (inset).

(legend continued on next page)

using flow cytometry [11, 14]. In cells that had not been previously exposed to galactose (i.e., naive cells), Gal1-mCherry was undetectable for the first 4 hr after shifting from glucose to galactose (Figures 1B and 1D, ACT). Between 4 hr and 8 hr after switching cells to galactose, expression of Gal1-mCherry was apparent in a subset of cells in the population (i.e., bimodal expression; Figures 1B and 1D, ACT). After 10 hr in galactose, the entire population expressed Gal1-mCherry (Figure 1D, ACT). In contrast, in cells that were previously grown in galactose and then repressed for 12 hr (~7 or 8 cell divisions; i.e., memory), the population responded uniformly and rapidly; Gal1-mCherry fluorescence was measurable within 4 hr after shifting back to galactose (unimodal expression; Figures 1B and 1D, REACT). Likewise, ectopic expression of *GAL1* promoted both rapid and unimodal accumulation of Gal1-mCherry (Figures 1B and 1D, ACT + *eGAL1*) [11, 14, 17, 24]. Furthermore, consistent with the model in Figure 1A, either loss of Gal80 or a point mutation in Gal4 (V864E) that disrupts the interaction with Gal80 [25] also resulted in rapid, unimodal expression of Gal1-mCherry (Figures 1C and 1D). Thus, memory leads to faster and more uniform *GAL1* transcriptional activation, most likely by promoting rapid and uniform relief of Gal80 repression.

To quantify the adaptive effect of faster reactivation of *GAL* genes during memory, we followed the growth kinetics upon shifting cells from glucose to galactose (Figure 1E). Naive cells exhibited a long growth lag before entering exponential phase (Figure 1E, ACT). In contrast, during memory or in cells ectopically expressing Gal1, adaptation was much faster (Figure 1E, REACT and ACT + *eGAL1*). Although the growth rates were ultimately similar once cells reached exponential phase, memory confers a large fitness benefit by decreasing the growth lag after shifting cells from glucose to galactose (Figure 1E, gray circles).

Rapid *GAL* gene activation in fungal species is also associated with increased responsiveness to low concentrations of galactose [26–28]. During memory or in cells expressing ectopic Gal1, Gal1-mCherry was expressed at higher levels in media with low concentrations of galactose (Figure 1F). Because yeast cells are most likely exposed to mixtures of sugars in nature, we asked whether this higher sensitivity for galactose also impacts the expression of Gal1-mCherry in the presence of glucose. *S. cerevisiae* normally does not induce *GAL* genes in the presence of low levels of glucose (0.2% glucose and 1.8% galactose; Figure 1G, inset). However, during memory or in the presence of ectopic Gal1, Gal1-mCherry expression was observed in the presence of glucose (Figure 1G, inset). This Gal1-mCherry expression correlated with a fitness benefit in 0.2% glucose + 1.8% galactose medium. In this medium, once glucose is exhausted after ~7 hr of growth, naive cells exhibited a significant lag before adapting to galactose (Figure 1G) [27, 29, 30]. However, during memory or in the presence of ectopic Gal1, this lag was absent and cells adapted immediately to galactose (Figure 1G). Thus, transcriptional memory provides a strong

adaptive advantage in both galactose and glucose-galactose mixtures.

Gal1-D117V Disrupts the Interaction with Gal80, Specifically Blocking *GAL* Transcriptional Memory

To explore the molecular basis of faster reactivation of *GAL* genes during memory, we performed a genetic screen based on fluorescence-activated cell sorting (FACS). After 4 hr in galactose, strong expression of Gal1-mCherry occurs during reactivation, but not during activation (Figures 1D and 2A). We exploited this difference to sort for *GAL* memory mutants: UV-mutagenized cells that failed to express Gal1-mCherry after 4 hr of reactivation (Figure 2A; sort I) but expressed Gal1-mCherry after 12 hr in galactose (Figure 2A; sort II). This second sort removed Gal[−] mutants or those that had lost Gal1-mCherry expression. The recovered cells were colony purified and screened by flow cytometry to identify those that specifically lost rapid *GAL1* reactivation during memory.

Based on the model in Figure 1A, we expected to identify null alleles of Gal1. We removed many such mutants by focusing on those that were able to grow on galactose (Gal1 is required to grow on galactose; data not shown). The screen also identified an allele of *GAL1* that was Gal⁺ but specifically blocked memory; *gal1-D117V* reduced the rate of Gal1-mCherry reactivation during memory without altering the rate of activation of Gal1-mCherry (Figure 2B). Reconstruction of the *gal1-D117V* mutation into the *GAL1* locus recapitulated these phenotypes (data not shown), confirming that this mutation is causative. *gal1-D117V* cells also lost the fitness benefit of memory; the growth of *gal1-D117V* during reactivation closely resembled the growth of naive wild-type cells during activation (Figure 2C). This mutation had no effect on Gal1-mCherry stability (Figure S1B) or the rate of activation (Figure 2B) and only slightly affected the rate of exponential growth in galactose (Figure 2C). Finally, the effects of this mutation were recessive because ectopic expression of Gal1 in *gal1-D117V* led to faster Gal1-mCherry expression (Figure S1C) and rapid adaptation to growth in galactose (Figure S1D). Thus, the *gal1-D117V* mutation specifically disrupts memory without significantly affecting other functions of Gal1.

The structures of Gal1, Gal3, and Gal3-Gal80 are known [20, 31]. Gal1 and Gal3 show 74% sequence identity and are structurally superimposable with a root mean square deviation of ~1.1 Å (Figure S1A) [20, 31]. Aspartate 117 maps to the predicted interaction surface between Gal1 and Gal80. In the Gal3-Gal80 structure, Gal3-Asp111 is at the structurally equivalent position to Gal1-Asp117 and forms an ionic bond with Gal80-Arg367 (Figures 2D and S1A) [20]. To test whether disrupting this salt bridge reduces the affinity between Gal1 and Gal80, we performed co-immunoprecipitation of wild-type and D117V Gal1-mCherry with Gal80-myc. Although these proteins were expressed at similar levels, immunoprecipitation

(E) Lysates from strains expressing Gal80-13xmyc and Gal1-mCherry were subjected to co-immunoprecipitation using anti-myc antibody. The immunoprecipitated fractions (IP) (top), the input fractions (middle), and the supernatant fraction after immunodepletion (bottom) were resolved by SDS-PAGE and immunoblotted against either mCherry (top two panels) or the myc epitope tag (bottom panels).

(F) Overlay of histograms for ACT and REACT of *gal1D117V* in (B).

The yeast strains and the number of biological replicates for all experiments are listed in Tables S1 and S2, respectively. Additional characterization of *gal1-D117V* mutant is shown in Figure S1.

of Gal80 recovered only ~20% of Gal1-D117V compared with wild-type Gal1 (Figure 2E). This reduced affinity for Gal80 led to slow, bimodal expression of Gal1-mCherry during both activation and reactivation (Figure 2F). Furthermore, a complementary mutant in Gal80 (R367L) predicted to disrupt the salt bridge between Gal80 and both Gal3 and Gal1 led to a Gal⁻ phenotype (data not shown). Thus, interaction between Gal1 and Gal80 plays a critical role in GAL gene transcriptional memory.

Constitutively Fast GAL Expression in *S. uvarum* Is due to Higher Basal Expression of Gal1

S. uvarum diverged from *S. cerevisiae* ~20 million years ago and has evolved a distinct strategy for adapting to growth in galactose (Figure S2A) [27, 32, 33]. We asked whether this species benefits from previous growth in galactose. Although the rate of Gal1-mCherry reactivation during memory was slightly faster than the rate of activation in *S. uvarum* (Figures 3A and S2B), this difference was much smaller than that observed in *S. cerevisiae* (Figure 3B). Moreover, in *S. uvarum*, previous growth in galactose did not lead to a fitness benefit (Figure 3C). This suggests that the rapid initial GAL gene induction in *S. uvarum* is sufficient to provide maximal fitness benefit and that increasing this rate further provides no additional effect.

Several differences between *S. uvarum* and *S. cerevisiae* might explain the difference in their response to previous growth in galactose; *S. uvarum* has higher basal expression of the key activators GAL1, GAL3, and GAL4 and lower expression of the GAL80 inhibitor [27, 33]. Thus, differences in *cis*-acting elements in the promoter, *trans*-acting factors, or both could lead to constitutive fast GAL gene expression. To investigate these possibilities, we substituted the GAL1 promoter (P_{GAL1}) in *S. cerevisiae* with P_{GAL1} from *S. uvarum*. In this strain, induction of Gal1-mCherry during both activation and reactivation was as fast as reactivation in wild-type cells (Figure 3D). Thus, P_{GAL1} from *S. uvarum* is sufficient to induce constitutively fast GAL1 expression in *S. cerevisiae* without any other *uvarum* factors.

The effects of P_{GAL1} from *S. uvarum* are consistent with this promoter being more easily induced. Hybrid *cerevisiae-uvarum* promoters suggest that this effect is largely explained by differences in the UAS_{GAL} elements and GAL1-proximal sequences (Figures S2C–S2E) [34]. However, because epigenetic GAL gene transcriptional memory in *S. cerevisiae* requires only a few hundred molecules of Gal1 per cell [14], very low basal expression of Gal1 by P_{GAL1} from *S. uvarum* might also cause this faster induction through a positive feedback mechanism. To distinguish between these possibilities, we asked whether *S. cerevisiae* with the *S. uvarum* P_{GAL1} also promoted faster activation of other GAL genes in *trans*. In cells bearing the *S. uvarum* P_{GAL1} , the rate of Gal7-Venus activation and reactivation was as fast as that observed during memory in wild-type *S. cerevisiae* (Figure 3E). Likewise, *S. uvarum* P_{GAL1} promoted faster adaptation to galactose (Figure 3F). Thus, the *S. uvarum* P_{GAL1} is sufficient to induce constitutive fast activation of GAL genes in *trans* and faster adaptation to galactose, most likely through basal Gal1 production.

Whereas the levels of basal Gal1 protein in cells having the P_{GAL1} from *S. uvarum* were below the level of detection using either flow cytometry or immunoblot, qRT-PCR revealed that

this promoter led to a significant increase in GAL1 mRNA in cells grown in glucose ($p = 0.03$; Student's *t* test; Figure 3F, inset) [27]. If this basal expression were causative, then disrupting the interaction between Gal1 and Gal80 should block this effect. Indeed, introduction of the *gal1-D117A* mutation into the *S. cerevisiae* strain harboring the *S. uvarum* P_{GAL1} blocked the *cis* and *trans* effects of this promoter on expression (Figures 3G and 3H) and growth (Figure 3I). Thus, constitutive fast GAL genes induction conferred by the *S. uvarum* P_{GAL1} is due to genetically encoded basal expression of Gal1 that impinges upon the same molecular mechanism employed during epigenetic transcriptional memory in *S. cerevisiae*.

Fitness Costs of Constitutive GAL1 Expression

If faster GAL genes expression promotes adaptation to galactose, why is it restricted to memory in *S. cerevisiae*? Basal Gal1 expression is detrimental to growth in glucose-galactose mixtures because the galactose-1-phosphate generated by the galactokinase activity of Gal1 inhibits phosphoglucosmutase and slows glycolysis [27, 35]. Consistent with this model, *S. cerevisiae* expressing ectopic GAL1 or *S. uvarum* showed a measurable growth disadvantage when adapting from glucose to a glucose-galactose mixture (1:1 ratio; 1% each sugar; Figures S3A and S3B). Competitive growth experiments between wild-type *S. cerevisiae* and *S. cerevisiae* harboring the *S. uvarum* P_{GAL1} also showed a fitness cost to basal expression of Gal1 (Figure S3C). Thus, consistent with previous studies, basal GAL1 expression can have both positive and negative effects on fitness; it promotes adaptation from glucose to galactose but is detrimental in glucose-galactose mixtures [29, 36].

The Gal4 CD Promotes Stronger Transcription during GAL Memory

In addition to the *gal1-D117V* mutant, which showed specific loss of memory without strong effects on activation, the flow cytometry screen also identified a mutation in Gal4 (*L282P*) that both blocked memory and led to defective activation of Gal1-mCherry (Figure S4A). This mutation most likely destabilizes the Gal4 protein, leading to lower protein levels (Figure S4A, inset). However, this mutation was interesting because it was resistant to ectopic expression of Gal1 (Figure S4A), confirming that the loss of memory in *gal4-L282P* cells was not simply an effect of lower levels of Gal1 during reactivation. Thus, Gal4-L282P both causes a defect in activation and blocks memory downstream of Gal1.

The *gal4-L282P* mutation lies within the CD of Gal4 (Gal4-239-767; Figure 4A) [37]. In other members of the zinc binuclear cluster transcription factor family, the CD has been proposed to have a regulatory function [38–41]. However, the role of Gal4 CD is unclear; deletion of this domain produces a largely functional activator, but mutations in this domain disrupt Gal4 function [37, 42].

To explore the role of the Gal4 CD, we tested how loss of this domain affected memory and the response to Gal1. Unlike Gal4-L282P, Gal4 Δ cd protein levels were similar to full-length Gal4 protein levels (Figure 4B, inset) and the rate of Gal1-mCherry activation was similar in *gal4 Δ cd* and wild-type cells (Figure 4B) [37]. However, cells lacking the CD showed no memory




Figure 3. Recently Diverged *Saccharomyces* Species Utilize Genetic and Epigenetic Strategies to Adapt to Growth in Galactose

Cells were shifted from glucose to galactose for activation (ACT) or grown in galactose overnight, repressed for 12 hr (*S. cerevisiae*) or 18 hr (*S. uvarum*) in glucose, and then shifted to galactose for reactivation (REACT).

(A) Gal1-mCherry fluorescence during activation and reactivation in *S. cerevisiae* and *S. uvarum*, normalized to expression at 10 hr.

(B) Ratio of reactivation to activation from the data in (B).

(C) OD₆₀₀ of *S. uvarum* during activation and reactivation.

(D–F) The *GAL1* promoter from *S. uvarum* was introduced in place of the endogenous *GAL1* promoter in *S. cerevisiae*. Gal1-mCherry (D) and Gal7-Venus (E) fluorescence relative to CFP and OD₆₀₀ (F) were measured during activation (ACT) and reactivation (REACT). Inset: basal *GAL1* mRNA, relative to *ACT1*, transcribed from the P_{GAL1} from *S. cerevisiae* and *S. uvarum* in glucose media.

(G–I) The *GAL1* promoter from *S. uvarum* driving expression of *GAL1* or *gal1-D117V* was introduced in place of the endogenous *GAL1* gene in *S. cerevisiae*. Gal1-mCherry (G) and Gal7-Venus (H) fluorescence relative to CFP and OD₆₀₀ (I) was measured during activation (ACT) and reactivation (REACT). Error bars represent SEM from ≥ 3 biological replicates for Gal1-mCherry fluorescence, ≥ 5 biological replicates for OD₆₀₀, and 12 replicates for RNA estimation. The yeast strains and the number of biological replicates for all experiments are listed in Tables S1 and S2, respectively. *GAL1* expression from P_{GAL1} from other *Saccharomyces* species is shown in Figure S2; growth trade-off for basal *GAL1* expression is shown in Figure S3.

(Figure 4B) and were unaffected by ectopic expression of either Gal1 (Figure S4B) or Gal3 (data not shown). Further, dimethylation of lysine 4 on histone H3 (H3K4me2), a chromatin mark that is associated with *GAL* transcriptional memory and is induced by ectopic expression of Gal1 [14], was still observed

in cells lacking the Gal4 CD (Figure S4D). Therefore, the *gal4 Δ cd* mutant blocked memory downstream of both Gal1 and chromatin changes associated with memory (Figure S4C).

Given the weak sequence conservation of the CD, we asked whether CD promotes memory by acting as a spacer to increase




Figure 4. The Gal4 CD Is Required for GAL Memory

(A) Schematic of the putative domain organization with a large CD of Gal4 (based on a structural prediction) between the N-terminal DNA binding domain and unstructured C-terminal activation domain.

(B–F) Naive cells (ACT), naive cells expressing ectopic GAL1 (ACT+eGAL1), or cells that were grown in galactose overnight and shifted to glucose for 12 hr (REACT) were shifted to galactose to assay the Gal1-mCherry fluorescence relative to constitutively expressed CFP.

(B) Wild-type and *gal4Δcd* mutant. Inset: immunoblot of Gal4-myc immunoprecipitated from wild-type and *gal4Δcd* mutant cells; arrows, Gal4; *, non-specific bands.

(C and D) CD of Gal4 was replaced with either 5 tandem repeats of β -spectrin domain (C) or the CD from Leu3 (D).

(E) Wild-type, *gal3Δ*, *gal4Δcd*, and *gal4Δcd gal3Δ* strains with or without eGAL1. Only the 0-hr and 10-hr time points are plotted for *gal3Δ* and *gal4Δcd gal3Δ* mutants.

(F) *gal4Δcd* strains with and without *gal80Δ* and *gal4V864E* mutation.

(G) Overlay of histograms of biological replicates from the indicated strains and time points in (B) and (F).

Error bars represent SEM from ≥ 3 biological replicates. The yeast strains and the number of biological replicates for all experiments are listed in Tables S1 and S2, respectively. Additional characterization of *gal4Δcd* mutant is shown in Figure S4.

the access of the activation domain to co-activators. The CD was replaced either with domains 12–16 of human β -spectrin, which should function as a spacer of similar size to the CD [43], or with the CD from Leu3, a related transcription factor [40, 44]. Although these hybrid proteins supported Gal1-mCherry expression, they blocked memory and were unresponsive to Gal1 (Figures 4C and 4D). Thus, the Gal4 CD has a sequence-specific

function in potentiating expression and is neither a simple spacer nor a generic, swappable domain.

Because Gal3 has a higher affinity for Gal80 than Gal1 [23], loss of memory could result if Gal4 Δ cd is de-repressed normally by Gal3 but is unresponsive to Gal1 (Figure 1A). To test this hypothesis, we asked whether Gal1 could replace Gal3 to promote activation of Gal1-mCherry. In cells lacking Gal3, Gal1-mCherry




Figure 5. An Inter-domain Interaction that Is Regulated by Gal80 Potentiates Gal4 Activation

(A) Confocal micrograph showing nuclear localization of the Gal4 CD-GFP expressed from P_{ADH1} (eCD-GFP) in cells having the nuclear envelope/endoplasmic reticulum marked with a RFP-tagged protein.

(B) Schematic for the experimental setup in (C)–(F). Gal80 dimer binds to the activation domain of Gal4 Δ cd and to eCD.

(C, D, and F) Gal1-mCherry fluorescence relative to CFP in presence of eCD and eCDmut (L282P).

(C) $gal4\Delta cd$ mutant with or without eGAL1. Inset: immunoblot of eCD-GFP and eCDmut-GFP. (D) $gal4\Delta cd$ or $gal4\Delta cd gal80\Delta$ mutants. (F) Wild-type cells with and without eGAL1.

(E) ChIP against eCD-GFP or eCDmut-GFP in the indicated strains. Recovery of the GAL1 promoter and a control locus, $PRM1$, were quantified relative to input by real-time qPCR. * $p \leq 0.05$ (Student's t test) relative to the ChIP enrichment of $PRM1$.

Error bars represent SEM from ≥ 3 biological replicates for Gal1-mCherry fluorescence and ≥ 3 replicated for ChIP experiments. The yeast strains and the number of biological replicates for all experiments are listed in Tables S1 and S2, respectively. Further characterization of eCD is shown in Figure S5. See also Figure S4.

($gal4$ -V864E) significantly increased the overall rate of activation in the $gal4\Delta cd$ cells (Figures 4F and 4G). Thus, the Gal4 CD operates downstream of Gal80 to regulate bimodal expression and potentiates maximal expression during memory.

If loss of the Gal4 CD completely blocked Gal80 repression, it should lead to expression of Gal1-mCherry in raffinose medium because Gal80 is the sole

repressor of GAL genes expression in raffinose. In raffinose, whereas either loss of Gal80 or loss of the interaction between Gal4 and Gal80 ($gal4$ -V864E) [25] led to de-repression of Gal1-mCherry, deletion of the CD alone did not (Figure S4C). Thus, loss of the Gal4 CD does not disrupt Gal80 repression but leads to faster relief of Gal80 repression.

The Gal4 CD Is a Target of Gal80 Repression

Loss of the Gal4 CD also altered Gal80 repression. During both activation and reactivation, $gal4\Delta cd$ cells showed unimodal Gal1-mCherry expression (Figures 4G and S4E). Hence, loss of the CD had two effects: it both reduced the strength of Gal1-mCherry expression during reactivation (as measured by average expression in the population; Figure 4B) and led to a more uniform activation of the population (Figures 4G and S4E). Because both loss of Gal80 and transcriptional memory also led to unimodal activation (Figure 1D), this implied that the CD is required for proper Gal80 repression. If so, then loss of Gal80 might not further increase the rate of activation. Indeed, neither loss of Gal80 nor disruption of the Gal4-Gal80 interaction

repressor of GAL genes expression in raffinose. In raffinose, whereas either loss of Gal80 or loss of the interaction between Gal4 and Gal80 ($gal4$ -V864E) [25] led to de-repression of Gal1-mCherry, deletion of the CD alone did not (Figure S4C). Thus, loss of the Gal4 CD does not disrupt Gal80 repression but leads to faster relief of Gal80 repression.

An Inter-domain Interaction Potentiates Gal4 Activation

In the other members of the Gal4 transcription factor family, the CD directly interacts with the activation domain to allosterically regulate activation [38, 40, 41]. To test whether the CD interacts with the rest of Gal4, we asked whether this domain could potentiate Gal4 Δ cd activation in *trans* (Figure 5B, schematic). Ectopically expressed CD localized in the nucleus (Figure 5A), independent of Gal4 (Figure S5A). Ectopic CD increased the rate of Gal1-mCherry activation in $gal4\Delta cd$ strains (Figures 5C and 5D). However, this effect required either expression of ectopic Gal1 (Figure 5C) or loss of Gal80 (Figure 5D). Chromatin immunoprecipitation (ChIP) experiments revealed that ectopic CD was recruited to the GAL1 promoter by Gal4 Δ cd (Figure 5E). This




Figure 6. Model for Epigenetic Potentiation of Gal4 Activation through Inter-domain Potentiation

(A) In wild-type cells during early activation, Gal80 repression is relieved in a subset of population, leading to lower level expression. Inter-domain interaction between CD and activation domain potentiates higher activation levels in cells relieved of Gal80.

(B) During memory (or in the presence of basal Gal1 expression), Gal80 repression is relieved early in whole population, leading to unimodal, fully potentiated *GAL* gene expression.

(C) *gal4Δcd* cells show uniform, low-level activation.

suggests that CD physically interacts with Gal4Δcd to potentiate activation, and this interaction is regulated by Gal1-Gal80.

This effect was highly specific; in the absence of ectopic Gal1, ectopic CD neither upregulated Gal1-mCherry expression nor bound to the *GAL1* promoter (Figures 5C, 5E, and S5B). Furthermore, ectopic CD neither bound nor potentiated activation from full-length Gal4 (Figures 5E, 5F, and S5C). Finally, ectopic L282P mutant CD (CDmut) (expressed at similar levels to wild-type CD; Figure 5C, inset) was not recruited to Gal4Δcd (Figure 5E). Thus, the L282P mutation in the Gal4 CD disrupts the interaction with other domains of Gal4, blocking potentiation.

DISCUSSION

This study provides important new insights into both the molecular mechanism of epigenetic *GAL* gene transcriptional memory in *S. cerevisiae* and an illustration of the evolutionary logic whereby the same molecules can produce either conditional,

epigenetic mechanisms of faster reactivation or constitutive, genetic mechanisms of fast activation. Our current model for both is shown in Figure 6. Gal80 physically interacts with both the activation domain and CD of Gal4, which may explain why ectopic CD is only able to complement the *gal4Δcd* phenotype in either the absence of Gal80 or in the presence of ectopic Gal1 [21, 45–48]. Although it is not yet clear whether the Gal80 that binds to the Gal4 activation domain is the same molecule as the Gal80 that interacts with the CD, our results suggest that both interactions are required for proper repression. Early during activation, Gal3 interacts with Gal80, permitting Gal4-mediated transcriptional activation in a subset of the cells in the population (Figure 6A). In these cells, the CD potentiates activation, leading to high-level expression. During memory, or in *S. uvarum*, the population shows uniform, rapid transition to high-level expression of *GAL* genes because of elevated concentrations of the Gal1 co-activator (Figure 6B). However, in cells lacking the Gal4 CD, the population responds uniformly, but the

level of expression is low. These cells do not show memory both because they are less well repressed by Gal80 (and therefore do not benefit from previous expression of Gal1) and because they are unable to achieve full activation.

Slight differences in the degree of repression of *GAL1* among *Saccharomyces* species leads to two different strategies that favor growth under different conditions. Low-level basal *GAL1* expression in *S. uvarum* leads to rapid adaptation to galactose but also encumbers a fitness cost in glucose-galactose mixtures [27, 29, 36]. On the other hand, tight *GAL1* repression restricts fitness during initial induction in galactose but leads to optimal utilization of glucose in the presence of galactose. Glucose is the most efficiently utilized sugar through glycolysis, and *S. cerevisiae* has a clear preference for it; expression of several genes is optimized for growth in glucose over other carbon sources [49]. Epigenetic memory in *S. cerevisiae* allows cells to benefit from recent growth in galactose without compromising the preference for glucose over longer timescales.

A whole-genome duplication during *Saccharomyces* evolution has led to specialization of function between the duplicated paralogs Gal1 and Gal3 [26, 28, 32–34]. The sub-functionalization of these proteins in different species has led to different evolutionary paths. *K. lactis* and *C. albicans*, which diverged from *Saccharomyces* before the whole-genome duplication, also exhibit constitutive fast induction of *GAL* genes [26, 28, 32]. This is because these species lack Gal3 and therefore must express higher basal levels of Gal1 to allow expression of the *GAL* genes. This implies that basal *GAL1* expression is the ancestral regulatory scheme that has been maintained in *S. uvarum*, in part through promoter-driven basal expression of Gal1 [26, 34, 50]. Replacing P_{GAL1} in *S. cerevisiae* with the P_{GAL1} from the more closely related *Saccharomyces* species *S. mikatae* and *S. paradoxus* did not lead to faster induction of Gal1 (Figures S2A and S2B). This suggests that basal *GAL1* expression due to promoter differences persisted in *S. uvarum* but was lost in *S. cerevisiae*, *S. paradoxus*, and *S. mikatae*. Tighter *GAL1* repression has been accompanied by specialization of *GAL3* as a co-activator: Gal3 from *S. cerevisiae* has lost galactokinase activity and has 10-fold higher affinity for Gal80 repressor than Gal1 [21, 23, 27, 33, 34]. Taken together, our results suggest that *GAL* transcriptional memory in *S. cerevisiae* is a product of tighter *GAL1* repression and specialization of *GAL3* as a co-activator. Thus, *GAL* memory may be an example where an epigenetic mechanism for faster reactivation evolved from an ancestral state of genetically encoded fast activation.

Using a FACS-based genetic screen, we identified two mutations that provide important insight into the molecular mechanism of *GAL* transcriptional memory. The *gal1-D117V* mutation maintains galactokinase function but reduces affinity for Gal80, specifically disrupting memory. Furthermore, *gal1-D117V* blocked fast *GAL* gene activation caused by the *S. uvarum* P_{GAL1} , confirming that these effects are mediated by low-level expression of Gal1.

The screen also identified *gal4-L282P*, a mutation in the CD of Gal4 that blocks the ability of Gal4 to respond to Gal1. Deletion of the CD disrupted memory without strongly altering Gal4 protein levels or the rate of activation. The CD has two functions: it promotes tighter Gal80 repression and it promotes stronger Gal4 activity (Figure 6C). Disrupting these functions resulted in a

qualitative change in the *GAL1* transcriptional output, leading to a more uniform population of cells that transitioned to a weaker level of expression (Figure 6C). In other words, unimodal induction is necessary, but not sufficient, for the rapid expression observed during memory. Because Gal80 interacts with both the CD and the activation domain [21, 45–48], we propose that the CD either enhances Gal80 recruitment to Gal4 or inhibits dissociation of Gal80 from Gal4. If so, then Gal80 would most likely also regulate the potentiation of Gal4 activation by the CD. We envision that the CD physically interacts with the activation domain, allosterically altering its ability to promote transcription. Consistent with this notion, the CD can interact with Gal4 Δ cd in *trans* to increase the rate of activation, and this interaction is both regulated by Gal80 and disrupted by substitution of Proline for Leucine 282. Thus, the CD plays a critical role in promoting both stronger repression by Gal80 and stronger transcription.

GAL transcriptional memory is a manifestation of ongoing resolution of expression levels of the partially redundant paralogs, Gal1 and Gal3. Among different *Saccharomyces* species, the degree of repression of *GAL1* in glucose dictates whether faster adaptation to galactose is a hardwired, genetic mechanism or a conditional, epigenetic mechanism. Because basal *GAL1* expression compromises fitness in mixtures of sugars, *S. cerevisiae* has traded faster kinetics of *GAL* gene activation for optimal growth in glucose-galactose mixtures. But during memory, *S. cerevisiae* can more rapidly adapt to a challenge that they have experienced recently by switching from a heterogeneous to uniform behavior and employing an inter-domain potentiation of Gal4 activation.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- METHOD DETAILS
 - Plasmids, yeast strains, and molecular biology
 - Flow cytometry
 - Genetic Screen
 - Growth Assay
 - Chromatin Immunoprecipitation
 - Microscopy
 - Competitive growth assay
- QUANTITATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental Information includes five figures and two tables and can be found with this article online at <https://doi.org/10.1016/j.cub.2017.10.035>.

AUTHOR CONTRIBUTIONS

J.H.B. and V.S. conceived the project, and V.S. carried out the experiments.

ACKNOWLEDGMENTS

The authors thank members of the Brickner laboratory and Professors Richard Gaber and Erik Andersen for helpful comments on the manuscript, Professor

Todd Hittenger (University of Wisconsin, Madison) for sharing *S. uvarum* strains, Meihua Christina Kuang for help with the growth assay, and Professor Mark Ptashne (Memorial Sloan Kettering Cancer Institute) for a fruitful discussion. This work was supported by NIH grant R01 GM118712 and American Heart Association pre-doctoral fellowship 14PRE18700074 (V.S.).

Received: July 19, 2017

Revised: September 18, 2017

Accepted: October 12, 2017

Published: November 16, 2017

REFERENCES

- López-Maury, L., Marguerat, S., and Bähler, J. (2008). Tuning gene expression to changing environments: from rapid responses to evolutionary adaptation. *Nat. Rev. Genet.* **9**, 583–593.
- Brickner, D.G., Cajigas, I., Fondufe-Mittendorf, Y., Ahmed, S., Lee, P.C., Widom, J., and Brickner, J.H. (2007). H2A.Z-mediated localization of genes at the nuclear periphery confers epigenetic memory of previous transcriptional state. *PLoS Biol.* **5**, e81.
- D'Urso, A., Takahashi, Y.H., Xiong, B., Marone, J., Coukos, R., Randise-Hinchliff, C., Wang, J.P., Shilatifard, A., and Brickner, J.H. (2016). Set1/COMPASS and Mediator are repurposed to promote epigenetic transcriptional memory. *eLife* **5**, e16691.
- Ding, Y., Fromm, M., and Avramova, Z. (2012). Multiple exposures to drought 'train' transcriptional responses in *Arabidopsis*. *Nat. Commun.* **3**, 740.
- Gialitakis, M., Arampatzis, P., Makatounakis, T., and Papamatheakis, J. (2010). Gamma interferon-dependent transcriptional memory via relocalization of a gene locus to PML nuclear bodies. *Mol. Cell. Biol.* **30**, 2046–2056.
- Guan, Q., Haroon, S., Bravo, D.G., Will, J.L., and Gasch, A.P. (2012). Cellular memory of acquired stress resistance in *Saccharomyces cerevisiae*. *Genetics* **192**, 495–505.
- Light, W.H., Brickner, D.G., Brand, V.R., and Brickner, J.H. (2010). Interaction of a DNA zip code with the nuclear pore complex promotes H2A.Z incorporation and INO1 transcriptional memory. *Mol. Cell* **40**, 112–125.
- Light, W.H., Freaney, J., Sood, V., Thompson, A., D'Urso, A., Horvath, C.M., and Brickner, J.H. (2013). A conserved role for human Nup98 in altering chromatin structure and promoting epigenetic transcriptional memory. *PLoS Biol.* **11**, e1001524.
- Seong, K.H., Li, D., Shimizu, H., Nakamura, R., and Ishii, S. (2011). Inheritance of stress-induced, ATF-2-dependent epigenetic change. *Cell* **145**, 1049–1061.
- Sung, S., He, Y., Eshoo, T.W., Tamada, Y., Johnson, L., Nakahigashi, K., Goto, K., Jacobsen, S.E., and Amasino, R.M. (2006). Epigenetic maintenance of the vernalized state in *Arabidopsis thaliana* requires LIKE HETEROCHROMATIN PROTEIN 1. *Nat. Genet.* **38**, 706–710.
- Zacharioudakis, I., Gligoris, T., and Tzamaras, D. (2007). A yeast catabolic enzyme controls transcriptional memory. *Curr. Biol.* **17**, 2041–2046.
- Acar, M., Becskei, A., and van Oudenaarden, A. (2005). Enhancement of cellular memory by reducing stochastic transitions. *Nature* **435**, 228–232.
- Pascual-Garcia, P., Debo, B., Aleman, J.R., Talamas, J.A., Lan, Y., Nguyen, N.H., Won, K.J., and Capelson, M. (2017). Metazoan nuclear pores provide a scaffold for poised genes and mediate induced enhancer-promoter contacts. *Mol. Cell* **66**, 63–76.e6.
- Sood, V., Cajigas, I., D'Urso, A., Light, W.H., and Brickner, J.H. (2017). Epigenetic transcriptional memory of GAL genes depends on growth in glucose and the Tup1 transcription factor in *Saccharomyces cerevisiae*. *Genetics* **206**, 1895–1907.
- Lohr, D., Venkov, P., and Zlatanova, J. (1995). Transcriptional regulation in the yeast GAL gene family: a complex genetic network. *FASEB J.* **9**, 777–787.
- Traven, A., Jelcic, B., and Sopta, M. (2006). Yeast Gal4: a transcriptional paradigm revisited. *EMBO Rep.* **7**, 496–499.
- Kundu, S., and Peterson, C.L. (2010). Dominant role for signal transduction in the transcriptional memory of yeast GAL genes. *Mol. Cell. Biol.* **30**, 2330–2340.
- Biggar, S.R., and Crabtree, G.R. (2001). Cell signaling can direct either binary or graded transcriptional responses. *EMBO J.* **20**, 3167–3176.
- Egriboz, O., Goswami, S., Tao, X., Dotts, K., Schaeffer, C., Pilauri, V., and Hopper, J.E. (2013). Self-association of the Gal4 inhibitor protein Gal80 is impaired by Gal3: evidence for a new mechanism in the GAL gene switch. *Mol. Cell. Biol.* **33**, 3667–3674.
- Lavy, T., Kumar, P.R., He, H., and Joshua-Tor, L. (2012). The Gal3p transducer of the GAL regulon interacts with the Gal80p repressor in its ligand-induced closed conformation. *Genes Dev.* **26**, 294–303.
- Platt, A., and Reece, R.J. (1998). The yeast galactose genetic switch is mediated by the formation of a Gal4p-Gal80p-Gal3p complex. *EMBO J.* **17**, 4086–4091.
- Timson, D.J., Ross, H.C., and Reece, R.J. (2002). Gal3p and Gal1p interact with the transcriptional repressor Gal80p to form a complex of 1:1 stoichiometry. *Biochem. J.* **363**, 515–520.
- Lavy, T., Yanagida, H., and Tawfik, D.S. (2016). Gal3 binds Gal80 tighter than Gal1 indicating adaptive protein changes following duplication. *Mol. Biol. Evol.* **33**, 472–477.
- Bhat, P.J., and Hopper, J.E. (1992). Overproduction of the GAL1 or GAL3 protein causes galactose-independent activation of the GAL4 protein: evidence for a new model of induction for the yeast GAL/MEL regulon. *Mol. Cell. Biol.* **12**, 2701–2707.
- Salmeron, J.M., Jr., Leuther, K.K., and Johnston, S.A. (1990). GAL4 mutations that separate the transcriptional activation and GAL80-interactive functions of the yeast GAL4 protein. *Genetics* **125**, 21–27.
- Dalal, C.K., Zuleta, I.A., Mitchell, K.F., Andes, D.R., El-Samad, H., and Johnson, A.D. (2016). Transcriptional rewiring over evolutionary time-scales changes quantitative and qualitative properties of gene expression. *eLife* **5**, e18981.
- Roop, J.I., Chang, K.C., and Brem, R.B. (2016). Polygenic evolution of a sugar specialization trade-off in yeast. *Nature* **530**, 336–339.
- Rubio-Teixeira, M. (2005). A comparative analysis of the GAL genetic switch between not-so-distant cousins: *Saccharomyces cerevisiae* versus *Kluyveromyces lactis*. *FEMS Yeast Res.* **5**, 1115–1128.
- Wang, J., Atolia, E., Hua, B., Savir, Y., Escalante-Chong, R., and Springer, M. (2015). Natural variation in preparation for nutrient depletion reveals a cost-benefit tradeoff. *PLoS Biol.* **13**, e1002041.
- Lee, K.B., Wang, J., Palme, J., Escalante-Chong, R., Hua, B., and Springer, M. (2017). Polymorphisms in the yeast galactose sensor underlie a natural continuum of nutrient-decision phenotypes. *PLoS Genet.* **13**, e1006766.
- Thoden, J.B., Sellick, C.A., Timson, D.J., Reece, R.J., and Holden, H.M. (2005). Molecular structure of *Saccharomyces cerevisiae* Gal1p, a bifunctional galactokinase and transcriptional inducer. *J. Biol. Chem.* **280**, 36905–36911.
- Kellis, M., Birren, B.W., and Lander, E.S. (2004). Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature* **428**, 617–624.
- Kuang, M.C., Hutchins, P.D., Russell, J.D., Coon, J.J., and Hittinger, C.T. (2016). Ongoing resolution of duplicate gene functions shapes the diversification of a metabolic network. *eLife* **5**, e19027.
- Hittinger, C.T., and Carroll, S.B. (2007). Gene duplication and the adaptive evolution of a classic genetic switch. *Nature* **449**, 677–681.
- de Jongh, W.A., Bro, C., Ostergaard, S., Regenber, B., Olsson, L., and Nielsen, J. (2008). The roles of galactitol, galactose-1-phosphate, and phosphoglucomutase in galactose-induced toxicity in *Saccharomyces cerevisiae*. *Biotechnol. Bioeng.* **101**, 317–326.

36. Venturelli, O.S., Zuleta, I., Murray, R.M., and El-Samad, H. (2015). Population diversification in a yeast metabolic program promotes antici-pation of environmental shifts. *PLoS Biol.* *13*, e1002042.
37. Ma, J., and Ptashne, M. (1987). Deletion analysis of GAL4 defines two tran-scriptional activating segments. *Cell* *48*, 847–853.
38. Des Etages, S.A., Saxena, D., Huang, H.L., Falvey, D.A., Barber, D., and Brandriss, M.C. (2001). Conformational changes play a role in regulating the activity of the proline utilization pathway-specific regulator in *Saccharomyces cerevisiae*. *Mol. Microbiol.* *40*, 890–899.
39. Rohde, J.R., Trinh, J., and Sadowski, I. (2000). Multiple signals regulate GAL transcription in yeast. *Mol. Cell. Biol.* *20*, 3880–3886.
40. Wang, D., Hu, Y., Zheng, F., Zhou, K., and Kohlhaw, G.B. (1997). Evidence that intramolecular interactions are involved in masking the activation domain of transcriptional activator Leu3p. *J. Biol. Chem.* *272*, 19383–19392.
41. Zhang, L., and Guarente, L. (1995). Heme binds to a short sequence that serves a regulatory function in diverse proteins. *EMBO J.* *14*, 313–320.
42. Hirst, M., Kobar, M.S., Kuriakose, N., Greenblatt, J., and Sadowski, I. (1999). GAL4 is regulated by the RNA polymerase II holoenzyme-associated cyclin-dependent protein kinase SRB10/CDK8. *Mol. Cell* *3*, 673–678.
43. Bhattacharyya, S., Renn, J.P., Yu, H., Marko, J.F., and Matouschek, A. (2016). An assay for 26S proteasome activity based on fluorescence anisotropy measurements of dye-labeled protein substrates. *Anal. Biochem.* *509*, 50–59.
44. Zhou, K.M., and Kohlhaw, G.B. (1990). Transcriptional activator LEU3 of yeast. Mapping of the transcriptional activation function and significance of activation domain tryptophans. *J. Biol. Chem.* *265*, 17409–17412.
45. Abramczyk, D., Holden, S., Page, C.J., and Reece, R.J. (2012). Interplay of a ligand sensor and an enzyme in controlling expression of the *Saccharomyces cerevisiae* GAL genes. *Eukaryot. Cell* *11*, 334–342.
46. Bhaumik, S.R., Raha, T., Aiello, D.P., and Green, M.R. (2004). In vivo target of a transcriptional activator revealed by fluorescence resonance energy transfer. *Genes Dev.* *18*, 333–343.
47. Leuther, K.K., and Johnston, S.A. (1992). Nondissociation of GAL4 and GAL80 in vivo after galactose induction. *Science* *256*, 1333–1335.
48. Sil, A.K., Alam, S., Xin, P., Ma, L., Morgan, M., Lebo, C.M., Woods, M.P., and Hopper, J.E. (1999). The Gal3p-Gal80p-Gal4p transcription switch of yeast: Gal3p destabilizes the Gal80p-Gal4p complex in response to galac-tose and ATP. *Mol. Cell. Biol.* *19*, 7828–7840.
49. Keren, L., Hausser, J., Lotan-Pompan, M., Vainberg Slutskin, I., Alisar, H., Kaminski, S., Weinberger, A., Alon, U., Milo, R., and Segal, E. (2016). Massively parallel interrogation of the effects of gene expression levels on fitness. *Cell* *166*, 1282–1294.e18.
50. Meyer, J., Walker-Jonah, A., and Hollenberg, C.P. (1991). Galactokinase encoded by GAL1 is a bifunctional protein required for induction of the GAL genes in *Kluyveromyces lactis* and is able to suppress the gal3 phenotype in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* *11*, 5454–5461.
51. Brickner, J.H., and Fuller, R.S. (1997). SOI1 encodes a novel, conserved protein that promotes TGN-endosomal cycling of Kex2p and other mem-brane proteins by modulating the function of two TGN localization signals. *J. Cell Biol.* *139*, 23–36.
52. Scannell, D.R., Zill, O.A., Rokas, A., Payen, C., Dunham, M.J., Eisen, M.B., Rine, J., Johnston, M., and Hittinger, C.T. (2011). The awesome power of yeast evolutionary genetics: new genome sequences and strain resources for the *Saccharomyces sensu stricto* genus. *G3 (Bethesda)* *1*, 11–25.
53. Sikorski, R.S., and Hieter, P. (1989). A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* *122*, 19–27.
54. Longtine, M.S., McKenzie, A., 3rd, Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., and Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* *14*, 953–961.
55. Brickner, D.G., Sood, V., Tutucci, E., Coukos, R., Viets, K., Singer, R.H., and Brickner, J.H. (2016). Subnuclear positioning and interchromosomal clustering of the GAL1-10 locus are controlled by separable, interdepen-dent mechanisms. *Mol. Biol. Cell* *27*, 2980–2993.
56. Oftedal, P. (1968). A theoretical study of mutant yield and cell killing after treatment of heterogeneous cell populations. *Hereditas* *60*, 177–210.
57. Eckardt, F., and Haynes, R.H. (1977). Kinetics of mutation induction by ultraviolet light in excision-deficient yeast. *Genetics* *85*, 225–247.
58. Ahmed, S., Brickner, D.G., Light, W.H., Cajigas, I., McDonough, M., Froysheter, A.B., Volpe, T., and Brickner, J.H. (2010). DNA zip codes con-trol an ancient mechanism for gene targeting to the nuclear periphery. *Nat. Cell Biol.* *12*, 111–118.
59. Brickner, J.H., and Walter, P. (2004). Gene recruitment of the activated INO1 locus to the nuclear membrane. *PLoS Biol.* *2*, e342.
60. Egecioglu, D.E., D’Urso, A., Brickner, D.G., Light, W.H., and Brickner, J.H. (2014). Approaches to studying subnuclear organization and gene-nuclear pore interactions. *Methods Cell Biol.* *122*, 463–485.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-GFP	AbCam (Cambridge, MA)	RRID: AB_3033395
Anti-H3K4me2	AbCam (Cambridge, MA)	RRID: AB_732924
Oligonucleotides		
GAL NB F: 5'-CCCCACAAACCTTCAAATTAACG-3'	IDT (Skokie, IL)	N/A
GAL NB R: 5'-CGCTTCGCTGATTAATTACCCC-3'	IDT (Skokie, IL)	N/A
PRM1CDS F: 5'-TTAGTCTTTGGGTCAATGTTCTCTG-3'	IDT (Skokie, IL)	N/A
PRM1CDS R: 5'-ATCAGCAGTGCTTTCAAACATGGAA-3'	IDT (Skokie, IL)	N/A
ACT1CDS F: 5'-GGTTATTGATAACGGTCTGGTATG-3'	IDT (Skokie, IL)	N/A
ACT1CDS R: 5'-ATACCTTGGTGTCTTGGTCTAC-3'	IDT (Skokie, IL)	N/A
Chemicals, Peptides, and Recombinant Proteins		
Bactopectone	VWR International (Pittsburgh, PA)	Cat#61001-506
Granulated Yeast Extract	VWR International (Pittsburgh, PA)	Cat#EM1.03753.0500
Galactose	Sigma (St. Louis, MO)	Cat#G0750
Yeast Nitrogen Base	Sunrise Science Products (San Diego, CA)	Cat#1500-250
CSM+Ade	Sunrise Science Products (San Diego, CA)	Cat#1128-100
Restriction enzymes	New England Biolabs (Ipswich, MA)	N/A
Dynabeads M-280 sheep anti-rabbit IgG	ThermoFisher	Cat#11203D
Experimental Models: Organisms/Strains		
Table S1	N/A	N/A

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Jason Brickner (j-brickner@northwestern.edu)

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All *S. cerevisiae* strains (W303 background) were generated from either CRY1 (*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1*) or CRY2 (*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1*; [51]) parent strain. *S. uvarum* strains were generated from JRY8153 strain from the Hittinger lab [52]. Yeast strains used in this study appear in Table S1. For flow cytometric estimation of *GAL1* expression, Gal1 was C-terminally tagged with mCherry. A constitutively expressed CFP or Venus was used as an internal reference.

METHOD DETAILS

Plasmids, yeast strains, and molecular biology

Plasmids for constitutive expression of *GAL1*, *GAL3* and CD of Gal4 (*CD*) were generated by amplifying these inserts with appropriate restriction site overhangs and cloning downstream of *ADH1* promoter (P_{ADH}) into pRS304 and pRS306 [53]. The resulting plasmids were linearized for insertion at *TRP1* and *URA3* loci, respectively. CFP and Venus fluorophores for normalization of *GAL1-mCherry* expression were cloned downstream of P_{TDH} in pRS306 vector and integrated at the *URA3* locus. Gene deletion and C-terminal tagging was performed as described previously [54]. Briefly, primers amplifying the deletion cassettes or fluorescent tags were guided by overhangs homologous to the target site and transformants were selected on the appropriate selection media. *GAL1*, *GAL4* and promoter mutants were generated by first inserting a *URA3-SUP4-o* cassette into the coding sequence, followed by transformation with a mutant PCR product and selection of FOA, as described previously [55]. The hybrid promoters were generated by stitching together promoter fragments through overlapping PCR. Cells were grown in Synthetic Dextrose Complete (SDC), Synthetic Galactose Complete (SGC) or Synthetic Raffinose Complete (SRC) at 25°C for growth rate studies. For expression and ChIP experiments cells were grown in Yeast Peptone Glucose (YPD) or Yeast Peptone Galactose (YPG), with the exception of *gal80Δ*,

which was growth SDC prior to induction with rich galactose media. In all media, except in [Figure 1E](#), the final concentration of total sugar was 2%.

Flow cytometry

Cells having *GAL1* tagged with mCherry were shifted from YPD to YPG and maintained between 0.05 to 0.3 OD₆₀₀ throughout the induction. 1 mL of culture was harvested at different times of induction and the cells were frozen and stored in 10% glycerol at -80°C . Cells were thawed on ice and passed through a BD LSRIL flow-cytometer in the Robert Lurie Comprehensive Cancer Center Flow Cytometry Core Facility. mCherry, CFP and Venus were excited with 561nm, 405nm and 488nm lasers, respectively. For detecting mCherry emission a 600nm long pass dichroic mirror and 610/20nm band pass filter set, for CFP emission 505nm long pass dichroic mirror and 525/50 band pass filter set and for Venus emission 530/30 band pass filter set was used. Roughly 5000 cells were analyzed to obtain the average intensities. The constitutively expressed CFP ($P_{TDH}\text{-CFP}$) and Venus ($P_{TDH}\text{-VENUS}$) served as a normalization control for Gal1-mCherry fluorescence in *S. cerevisiae* and *S. uvarum* respectively; Gal1 fluorescence intensity was expressed as ratio of Gal1-mCherry to CFP or Venus. Biological replicates for all *GAL1-mCherry* induction were performed at least three times using independent cultures on different days. The data analysis did not necessitate either randomization or blinding at any stage.

Genetic Screen

Exponentially growing wild-type cells in SGC were mutagenized by exposure to 254nm ultraviolet (UV) light, using a hand-held lamp (UVGA-25, UVP Inc). 10 mL of cells at OD_{600nm} of 0.1 were placed in a 10cm Petri Dish. These cells were exposed to UV for 60sec from a distance of 15cm to the UV source, which killed 30% of the cells [56, 57]. The mutagenized cells were transferred to YPD, grown for 12 hr and then shifted to YPG for 4 hr. Fluorescence activated cell sorting for non-fluorescent cells was done using the BD FACSaria SORP 5 at the Northwestern Flow Cytometry Core Facility. Approximately two million cells were harvested in YPG. Cells were pelleted and then resuspended in fresh YPG for additional 8 hr. Sorting was then performed for cells that have wild-type Gal1-mCherry expression levels. Cells collected from the second sort were plated for single colonies on galactose plates. *GAL1-mCherry* activation and reactivation kinetics was individually assayed for each colony. Complementation with wild-type *GAL* genes was used for mapping mutations that lead to specific reactivation defects followed by subsequent Sanger sequencing of the mutant loci to identify the mutation.

Growth Assay

Exponentially growing cells were diluted to an OD₆₀₀ = 0.1, washed with media containing no sugar and then re-suspended in the media with the desired sugar. About 350 μL of each culture was added to a well in a 96-well flat bottom plate. Growth was monitored by measuring OD₆₀₀ every 20 min for 24 hr-40 hr using a 96-well plate reader (BioTek Synergy), normalized to media without cells. The cultures were not shaken at any time during the growth assay. The cell density at $t = 0$ was subtracted from all measurements. At least three biological replicates were done for each condition each time and repeated on different days.

Chromatin Immunoprecipitation

ChIP was performed as described previously with slight modifications [3, 8, 58, 59]. Approximately 100 mL culture of exponentially growing cells at OD_{600nm} of 0.8 was fixed with 1% formaldehyde for 15min at room temperature. Formaldehyde was quenched with a final concentration of 0.15M Glycine. Cells were then washed with 0.1M Tris pH 7.5 and collected through filtration. From this point up to elution, the cells were kept at $\sim 4^{\circ}\text{C}$. The cells were suspended in 600 μL cold lysis buffer (50mM HEPES pH 7.5, 140mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% Sodium deoxycholate) and vortexed with 600 μL of glass beads using Vortex Genie 2. The cells were vortexed five times at max speed for 1min, with 1min intervals. The lysed cells were harvested and spun at 7000 g for 10min. The supernatant was aspirated and the pellet containing the chromatin was re-suspended in 1 mL of lysis buffer. The re-suspended pellet was sonicated using a Branson Sonifier 450, for 15 pulses of 10sec, with 10sec intervals. The sonicated lysate was spun at 15000 g for 10min. The supernatant was harvested and diluted to $\sim 4\text{mg/mL}$ using lysis buffer. 50 μL of supernatant was taken as input control from each sample and mixed with 200 μL of 1%SDS. For immuno-precipitation, 8 μL of anti-Rabbit IgG Dynabeads were mixed with 2 μL of either anti-GFP or anti-H3K4me2, for each sample. The supernatant was incubated overnight at 4°C with the antibody bound beads. The beads were isolated using a magnetic separator and washed five times with 1 mL of lysis buffer. For elution of immuno-precipitated DNA, beads were incubated with 125 μL of elution buffer (50mM Tris pH 7, 10mM EDTA, 1% SDS) at 65°C for 15min. This was repeated twice and the two eluates were pooled. Both the input and the eluate were treated with 2 μL of 10mg/mL RNaseA at 37°C for 30min, followed by 2 μL of 20mg/mL of Proteinase K at 42°C for 2 hr. Un-crosslinking was done at 65°C for 8 hr and DNA fragments were eluted in 1X TE using QIAGEN PCR clean-up kit. The input and immuno-precipitated samples were diluted 400-fold and 10-fold, respectively. The recovery of the fragments from *GAL1*, *ACT1* and *PRM1* loci, by ChIP, were quantified by qPCR using primers specified in the key resource table. The enrichment was quantified as ratio of Immuno-precipitated DNA over input DNA (IP/Input). We excluded Ct values (qPCR) that were more than two cycles apart from the average for the same DNA samples.

Microscopy

The cells containing CD-GFP and ER/nuclear envelope targeted mCherry were grown in SDC overnight. The OD₆₀₀ was kept below 0.5 and a sample of this culture was directly spread on a slide for imaging. The SP5 Line Scanning Confocal Microscope (Leica Biosystems) at the Northwestern University Biological Imaging Facility was used for imaging as described earlier [60]. The Argon

488nm and Diode pumped solid-state 561nm lasers at ~10% power were used for exciting the CD-GFP and nuclear directed Cherry, respectively. The images were acquired for 150 μm X 150 μm field at 2048 X 2048 pixel resolution for ten z stacks of 0.73 μm with 0.34 μm step size through 100 X 1.44 NA objectives. The images from the GFP and Cherry channels were merged using the LAS AF Lite software from Leica.

Competitive growth assay

Two exponentially growing *S. cerevisiae* strains, in YPD, containing either *PGAL1uvrurum* or *PGAL1cerevisiae* were mixed together in equal ratio. The mixture was co-cultured in YP media containing 1% glucose and 1% galactose at 25°C for 36 hr with periodic dilutions. The OD_{600} for the cultures were kept below 0.5 at all times. Constitutively expressed Venus in one strain was used for flow-cytometric estimation of changes in relative fraction of the two populations over time. Parallel competitive assays were performed with Venus expressed in either one of the strain to control for the effect of Venus on growth. Malthusian fitness coefficients (M) were calculated as $M = \ln(10^{\log[(\text{GFP}_{\text{end}}/\text{BFP}_{\text{end}})/(\text{GFP}_{\text{start}}/\text{BFP}_{\text{start}})]/t})$ as described previously [34].

QUANTITATION AND STATISTICAL ANALYSIS

At least 3 biological replicates were performed for each experiment to plot the average and standard error of the means. The replicates were performed from cultures grown independently on different days. The data was plotted using the GG-plot. To evaluate the significance of difference between strains or treatments with respect to the reference, an unpaired, two-tailed Student's t test was performed. We did not perform additional tests for applicability of Student's t test to our data.